

THE STAR-FORMATION HISTORY OF THE UNIVERSE

MICHAEL ROWAN-ROBINSON

*Astrophysics Group, Blackett Laboratory,
Imperial College of Science Technology and Medicine
Prince Consort Road, London SW7 2BZ*

1. Introduction

Many authors have attempted to model the star formation history of the universe (eg Cole et al 1994, Pei and Fall 1995) or equivalently the evolution of the starburst galaxy population (eg Franceschini et al 1991, 1994, 1997, Blain and Longair 1993, Pearson and Rowan-Robinson 1996, Guiderdoni et al 1998, Dwek et al 1998, Blain et al 1999). Most of these studies explore a small range of models designed to fit the available observations. In the present paper I report the results of a parameterized approach to the star formation history of the universe, which allows a large category of possible histories to be explored and quantified. The parametrized models can be compared with a wide range of source-count and background data at far infrared and submillimetre wavelengths to narrow down the parameter space that the star formation history can occupy.

Throughout this paper I use the most recent Bruzual and Charlot star formation scenarios, with a Salpeter IMF with mass cutoffs at 0.1 and 100 $M_{\odot}$, for which the relevant conversion factors are:

$$L(H\alpha)/L_{\odot} = 10^{7.60} \dot{\phi}_*, \quad (1)$$

$$L(60\mu m)/L_{\odot} = 10^{9.66} \epsilon \dot{\phi}_*, \quad (2)$$

$$L(2800\text{\AA})/L_{\odot} = 10^{9.35} \dot{\phi}_*, \quad (3)$$

$$L(2000\text{\AA})/L_{\odot} = 10^{9.50} \dot{\phi}_*, \quad (4)$$

$$\dot{\phi}_* = 42 \dot{\phi}_Z, \quad (5)$$

where $\dot{\phi}_*$, $\dot{\phi}_Z$, are the star and metal formation rates in solar masses per year per cubic Mpc and ϵ is the fraction of opt-uv light absorbed by dust (Rowan-Robinson et al 1997). A Hubble constant of 50 $km/s/Mpc$ is used throughout, and $\Omega_o = 1$ (unless otherwise stated).

2. Star formation rate derived from ultraviolet data

I first estimate the star formation history at uv wavelengths, with no correction for the effects of dust. For the local, $z = 0$, star formation rate, I have gone back to the $H\alpha$ data of Gallego et al (1995) in order to remove the reddening correction they have applied. I find a mean reddening correction of 4.4, to give a value for $\dot{\phi}_*$ of $10^{-2.76 \pm 0.20} M_\odot/\text{yr}/Mpc^3$. I have also estimated the star-formation rate from a sample of 170 bright ($B \leq 13.8$) nearby ($V < 5000$ km/s), optically selected galaxies for which there are both $60 \mu\text{m}$ ($S(60) \geq 0.6$ Jy) and large-beam $2000 \text{ \AA}$ observations (Buat et al 1987, Kinney et al 1993, Deharveng et al 1994, Meurer et al 1995). The $60 \mu\text{m}$ and B-band luminosity functions can be used to verify that the sample is representative of all galaxies and allow the effective area surveyed to be estimated. The $2000 \text{ \AA}$ luminosity density is found to be $10^{18.42} W Mpc^{-3}$ and the corresponding star formation rate is $10^{-2.48 \pm 0.06} M_\odot \text{yr}^{-1} Mpc^{-3}$. This is consistent with the revised estimate given above from the Gallego et al $H\alpha$ data and with the estimate given by Lilly et al (1996) extrapolated from B-band data, $10^{-2.42} M_\odot \text{yr}^{-1} Mpc^{-3}$, but is a factor of 3 lower than the value used by Madau et al (1996).

Madau et al (1996) have given estimates for the star formation rate at $z = 2 - 4.5$ derived from analysis of Lyman drop-out galaxies in the Hubble Deep Field. This analysis appeared to demonstrate that the star formation drops off steeply at $z > 2$. Connolly et al (1997) have used photometric redshifts for the brighter HDF galaxies to derive the star formation rate at $z = 0.5 - 2$. Their method involves the use of J, H and K data in addition to the HST U,B,V,I data to determine the redshifts. Although this certainly greatly improves the reliability of the photometric redshifts, it restricts the analysis to $< 10\%$ of the galaxies detected in the HDF by HST.

To try to extend the results of Connolly et al to a wider range of redshift, and to test the robustness of the conclusions of Madau et al (1996), I have used the photometric redshift method of Mobasher et al (1996) to analyze a much larger sample of HDF galaxies, namely all those galaxies with $I_{AB} \leq 29.0$ from the catalogue of Williams et al (1996) detected in at least the I and V bands, a total of 2438 galaxies once duplicates have been eliminated. This method gives redshifts accurate to 13 % in $(1+z)$, although there is obviously a possibility of aliasing for galaxies detected in only 2 bands (22% of the sample). For these 2438 HDF galaxies I have estimated the luminosity functions of the galaxies at rest-frame $2800 \text{ \AA}$ in bins of 0.05 in $\log_{10}(1+z)$, for $z \leq 4$. The resulting star formation history is shown in Fig 1. Agreement with the results of Lilly et al (1996), Treyer et al (1997) and Connolly et al (1997) is excellent over the range $z = 0.2-2.5$. However the derived star formation rates at higher redshifts are considerably higher




Figure 1. Star formation history derived from ultraviolet data, with no correction for effects of dust. Data points are from Gallego et al (1995), (G), Treyer et al (1998) (T), Lilly et al (1996) (L), from the sample of bright local galaxies (section 2) at $z = 0.01$ (U), and from the analysis of photometric redshifts of HDF galaxies (open circles). The models shown are chosen to fit the CDM predictions of Cole et al (1996) $(P,Q) = (1.5, 5.)$, and to account for the far infrared and submm counts, $(P,Q) = (0.5, 5.5)$.

than those of Madau et al (1996) and shows no sign of a decline in star formation rate beyond $z = 1$.

3. Star formation history from infrared or submillimetre data, or from ultraviolet data with correction for the effects of dust

Figure 2 shows the star formation history derived either from ultraviolet or $H\alpha$ data with correction for reddening, or from far infrared or submillimetre data under the assumption that most ultraviolet and visible light from star forming regions is absorbed by dust (i.e. $\epsilon \simeq 1$).

In the case of the HDF data the extinction is derived using the grain properties of Rowan-Robinson (1992) assuming a (conservative) extinction corresponding to $E(B-V) = 0.1$ at the rest-frame wavelength corresponding to $\lambda_{obs} = 6000 \text{ \AA}$, which gives correction factors ranging from 1.5 at $z=0.2$ to 2.5 at $z = 3.8$. Most estimates of correction factors to be applied to high redshift star-forming galaxies range from 2-7 (Pettini et al 1997, Meurer et al 1998, Calzetti 1998, Steidel et al 1999). Note the good agreement of my values derived from the HDF with those of Steidel et al (1999) from




Figure 2. Star formation history derived from infrared (IRAS: Saunders et al 1990, ISO: Rowan-Robinson et al 1997, F: Flores et al 1997), submillimeter (SCUBA, 850 μ m: Hughes et al 1998), and from ultraviolet data with correction for effects of dust, from Gallego et al (1995) (Ga), Treyer et al (1998) (T), both corrected by a factor 2 for dust extinction, Gronwall (1998) (Gr) and Steidel et al (1999) (S) and from the analysis of photometric redshifts of HDF galaxies assuming $E(B-V) = 0.1$ (open circles).

surveys for Lyman break galaxies.

The ISO-HDF estimates of Rowan-Robinson et al (1997) for the redshift range 0.4-0.7 have been corrected for the fact that one of the 5 galaxies in this range is no longer thought to be a starburst (source 3 in Table 1 of Rowan-Robinson et al). This has an almost negligible effect on the total star formation rate. Also shown are the star formation rates derived by Hammer and Flores (1998) from ISO observations of a CFRS field.

The shape of the observed star formation history is similar to that derived from the uv (Fig 1), but the values are a factor 2-3 higher at most redshifts, implying $\epsilon = 2/3$ - $3/4$. For comparison Calzetti (1998) gives a range for ϵ of $1/2$ to $2/3$ for starburst galaxies.

4. Parametrized approach to star formation history

To study what constraints on the star formation history can be derived from source-counts and background intensity measurements at far infrared and submm wavelengths, I present here a parameterized approach to the problem, investigating a wide range of possible star formation histories.

The constraints we have on the star formation rate, $\dot{\phi}_*(t)$ are that: (i) it is zero for $t = 0$, (ii) it is finite at $t = t_o$, (iii) it increases with z out to at least $z = 1$ (and from (i) must eventually decrease at high z).

A simple mathematical form consistent with these constraints is

$$\dot{\phi}_*(t)/\dot{\phi}_*(t_o) = \exp Q(1-(t/t_o)) (t/t_o)^P \quad (6)$$

where P and Q are parameters ($P > 0$ to satisfy (i), $Q > 0$ to satisfy (iii)). I assume that $\dot{\phi}_*(t) = 0$ for $z > 10$.

Equation (6) provides a simple but versatile parameterization of the star formation history, capable of reproducing most physically realistic, single-population scenarios.

Given an assumed (P, Q) I then determine the $60 \mu\text{m}$ luminosity function, using the IRAS 1.2 Jy sample (Fisher et al 1995). I fit this with the form assumed by Saunders et al (1990)

$$\eta(L) = C_*(L/L_*)^{1-\alpha} e^{-0.5[\log_{10}(1+L/L_*)/\sigma]^2} \quad (7)$$

and find that the luminosity function parameters can be well approximated as follows:

$$\sigma = 0.765 - 0.04 W$$

$$\log_{10} L_* = 8.42 + 0.07 W - 2 \log_{10}(H_o/100)$$

where $W = 0.825 Q - P$, and I have assumed fixed values for $\alpha = 1.09$, $C_* = 0.027 (H_o/100)^3$. It is not clear that previous studies have correctly taken account of the need to change the $60 \mu\text{m}$ luminosity function as the rate of evolution is varied. The study of Guiderdoni et al (1998) explicitly violates the known constraints on the $60 \mu\text{m}$ luminosity function at the high luminosity end and as a result the models predict far too many high redshift galaxies at a flux-limit of 0.2 Jy at $60 \mu\text{m}$, where substantial redshift surveys have already taken place.

To transform this $60 \mu\text{m}$ luminosity function to other wavelengths I assume that the spectral energy distributions of galaxies are a mixture of two components, a starburst component and a 'cirrus' component (cf Rowan-Robinson and Crawford 1989). I have used the latest predictions for infrared seds of these two components by Efstathiou et al (1999).

There is no explicit allowance for emission from AGN in the models, though it is known that at low far infrared luminosities, about 10 % of galaxies contain AGN, with the proportion increasing towards higher luminosities. Provided the far infrared component peaking at $60 \mu\text{m}$ which I have identified as the 'starburst' component is indeed powered by a starburst (cf Rowan-Robinson and Crawford 1989, Rigopoulou et al 1996, Lutz et al 1998), it is irrelevant for estimates of the star formation rate whether the galaxy also contains an AGN. I am implicitly assuming that any correlation between far infrared luminosity and the optical/uv/X-ray luminosity from an accreting black hole is caused by a common feeding mechanism.

I can now predict the counts and background intensity at any wavelength and by comparing with observed values, constrain loci in the P-Q plane. Fig 3 shows a number of such loci for the case of pure luminosity evolution in an Einstein de Sitter model ($\Omega = 1$). The uncertainties on these loci are typically ± 0.5 in Q, except for the 60 μm counts, where the uncertainty is much broader (± 2 in Q), with rather weak dependence on P for $P > 0$. The locus shown for N60 is in fact the upper end of the $1-\sigma$ range for the number of sources per ster at 60 μm at 0.25 mJy. Figure 4,5 shows integral counts at 850 and 60 μm . Figure 6 shows the spectrum of the integrated background radiation, compared with selected models. Because of the simplistic nature of the assumed optical sed, I have not attempted to show optical counts, but it is of interest to try to get roughly the correct balance of optical and far infrared/submm background radiation.

We see that models consistent with the far infrared and submillimeter counts can be found, eg $(P,Q) = (0.5,5.5)$, and that such models also are consistent with the observed 850 μm background. However stronger evolution is required to fit the 140-350 μm background intensity, eg $(P,Q) = (0.5,6.5)$, and such models are inconsistent with the observed counts. The CDM model, which can be approximated by $(P,Q) = (1.5, 5.0)$, is consistent with the 60 μm counts but nothing else.

An important constraint on the models is that the total mass of stars produced in galaxies should be greater than or equal to the mass of stars observed, $\Omega_* \geq 0.006 \pm 0.0018(H_o/50)^{-1}$ (Lanzetta et al 1996), and that it should be less than the total density of baryons in the universe, $\Omega_* \leq 0.05 \pm 0.01(H_o/50)^{-2}$ (Walker et al 1991). The models which fit the counts ($(P,Q) = (0.5, 5.5)$) and 140-350 μm background ($(P,Q) = (0.5, 6.5)$), give $\Omega_* = 0.015$ and 0.032 respectively.

We might expect that the cosmological model could have a significant effect on the relationship between predicted counts and predicted background intensity, since the latter is sensitive to how the volume element and look-back time change with redshift.

To test this we have explored models with $\Lambda = 0$, for which all the required formulae are analytic. Figure 7 shows the P-Q diagram for a model with $\Omega_o = 0.3$, with loci similar to those of Fig 3 plotted. In this cosmological model there does appear to be a single star formation history which is consistent with both source-count and background radiation data, namely one with $P = 1.5$, $Q = 8.2$. Predicted counts and background spectrum are shown for this model as the dotted curve in Figs 4-6.

Thus within the framework of the types of star formation history considered in section 5, there is a clear sensitivity to cosmological parameters, with $\Omega_o = 0.3$ preferred to $\Omega_o = 1$. This not to say that more complex assumptions about the star formation history, for example the inclusion of




Figure 3. P-Q diagram, with loci for models fitting 60, 175 and 850 μm counts (solid curves) and 140, 240, 350 and 850 μm background intensity (dotted curves). Counts loci correspond to $N_{60}(0.25\text{Jy}) = 2.0$, $N_{175}(0.1\text{Jy}) = 39$ (Kawara et al 1998), $\log N_{850}(4\text{mJy}) = 900$ per sq deg (Hughes et al 1998). Background loci correspond to $\nu I_\nu(140) = 12.5$, $\nu I_\nu(240) = 11$, $\nu I_\nu(350) = 6.0$ and $\nu I_\nu(850) = 0.5 \text{ nW/m}^2/\text{sr}$ (Fixsen et al 1998).

a dust-enshrouded population not represented at low redshift as in Franceschini et al (1997), might not be consistent with $\Omega_o = 1$. The full dependence on cosmological parameters will be explored in later work.

References

- Blain A.W., Longair M.S., 1993, MN 264, 509
 Blain A.W., Smail I., Ivison R.J., Kneib J.-P., 1999, MN 302, 632
 Buat V., Donas J., Deharveng J.-M., 1987, AA 185, 33
 Calzetti F., 1998, in 'Dwarf Galaxies and Cosmology', ed. T.X.Thuan, C.Balkowski, V.Cayatte, J. Tran Thanh Van (Editions Frontieres) astro-ph/9806083
 Cole S., Aragon-Salamanca A., Frenk C.S., Navarro J.F., Zepf S.E., 1994, MN 271, 781
 Connolly A.Z., Szalay A.S., Dickinson M., Subbarao M.U., Brunner P.J., 1997, ApJ 486, L11
 Deharveng J.-M., Sasseen T.P., Buat V., Lampton M., Wu X., 1994, AA 289, 715
 Dwek E., et al, 1998, ApJ 508, 106
 Eales S., Lilly S.J., Gear W.K., Dunne L., Bond J.R., Hammer F., Le Fevre O., Crampton D., 1999, ApJL (in press), astro-ph/9808040
 Efstathiou A., Rowan-Robinson M., Siebenmorgen R., 1999, MN (in press)
 Fisher K.B., Huchra J.P., Strauss M.A., Davis M., Yahil A., Schlegel D., 1995, ApJS 100, 69
 Fixsen D.J., Dwek E., Mather J.C., Bennett C.L., Shafer R.A., 1998, ApJ 508, 123




Figure 4. Integral source counts at 850 μ m. Data are from Hughes et al (1998), Eales et al (1999), Smail et al (1997). The 3 models shown are, from bottom at faint fluxes, for $\Omega_o = 1$ and (P,Q) = (1.5, 5.0), (0.5, 5.5) and (0.5, 6.5) (solid curves) and for $\Omega_o = 0.3$, (P,Q) = (1.5, 8.2) (dotted curve).

- Franceschini A., Toffolatti L., Mazzei P., Danese L., De Zotti G., 1991, AAS 89, 285
 Franceschini A., Mazzei, P., de Zotti, G., Danese, L., 1994, ApJ 427, 140
 Franceschini A., Silva L., Fasano G., Granato G.L., Bressan A., Arnouts S., Danese L., 1998, ApJ 506, 600
 Gallego J., Zamorano J., Aragon-Salamanca A., Rego M., 1995, ApJ 455, L1
 Gregorich D.T., Neugebauer G., Soifer B.T., Gunn J.E., Herter T.L., 1995, AJ 110, 259
 Gronwall C., 1998, in 'Dwarf Galaxies and Cosmology', eds T.X.Thuan, C.Balkowski, V.Cayatte, J.Tran Thanh Van, (Editions Frontieres), astro-ph/9806240
 Guiderdoni B., Hivon E., Bouchet F.R., Maffei B., 1998, MN 295, 877
 Hacking P.B. and Houck J., 1987, ApJS 63, 311
 Hammer F., Flores H., 1998, in 'Dwarf Galaxies and Cosmology', eds T.X.Thuan, C.balkowski, V.Cayatte, J.Tran Thanh Van, (Editions Frontieres), astro-ph/9806184
 Hauser M.G., et al, 1998, ApJ 508,25
 Hughes D.H., et al, 1998, Nat 394, 241
 Kawara K., et al, 1998, AA (in press)
 Kinney A.L., Bohlin R.C., Calzetti D., Panagia N., Wyse R.F.G., 1993, ApJS 86, 5
 1989, MN 240, 349
 Lilly, S.J., Le Fevre, O., Hammer, F., Crampton, D., 1996, ApJ 460, L1
 Lonsdale C.J., Hacking P.B., Conrow T.P., Rowan-Robinson M., 1990, ApJ 358, 60
 Lutz D., Spoon H.W.W., Rigopoulou D., Moorwood A.F.M., Genzel R., 1998, ApJ 505, L103
 Madau, P., Ferguson, H.C., Dickinson, M.E., Giavalisco, M., Steidel, C.C., Fruchter, A., 1996, MNRAS 283, 1388
 Meurer G.R., Heckman T.M., Leitherer C., Kinney A., Robert C., Garnett D.R., 1995, AJ 110, 2665




Figure 5. Source counts at $60\ \mu\text{m}$. Data are from Lonsdale et al (1990) (at 0.2-10 Jy), Hacking and Houck (1987) (at 50-100 mJy), Gregorich et al (1995) (higher point at 50 mJy). Models as in Fig 4.




Figure 6. Predicted spectrum of integrated background for same models as Fig 4. Data from Fixsen et al (1998) (far ir and submm), Pozzetti et al (1998) (opt and uv).




Figure 7. P-Q diagram for $\Omega_o = 0.3$ model (section 7), with loci for models fitting 60, 175 and 850 μm counts (solid curves) and 140 and 850 μm background intensity (dotted curves).

- Mobasher, B., Rowan-Robinson, M., Georgakakis, A., Eaton, N., 1996, MN 282, L7
 Pearson, C., Rowan-Robinson, M., 1996, MN 283, 174
 Pettini M., et al, 1997, in 'Cosmic Origins: Evolution of Galaxies, Stars, Planets and Life', eds Woodward C.E., et al, ASP Conf.Ser. (astro-ph/9708117)
 Pei, Y.C., and Fall, S.M., 1995, ApJ 454, 69
 Pozzetti L., Madau P., Ferguson H.C., Zamorani G., Bruzual G.A., 1998, MN 298, 1133
 Puget, J.-L., Abergel A., Bernard J.-P., Boulanger F., Burton W.B., Desert F.-X., Hartmann D., 1966, AA 308, 5
 Rigopoulou D., Lawrence A., Rowan-Robinson M., 1996, MN 288, 1049
 Rowan-Robinson M., and Crawford J., 1989, MN 238, 523
 Rowan-Robinson, M., 1992, MN 258, 787
 Rowan-Robinson M. et al, 1997, MN 289, 490
 Saunders, W., Rowan-Robinson, M., Lawrence, A., Efstathiou, G., Kaiser, N., Frenk, C.S., 1990, MN 242, 318
 Schlegel D.J., Finkbeiner D.P., Davis M., 1998, ApJ 500, 525
 Smail I., Ivison R.J., Blain A.W., 1997, ApJL 490, L5
 Steidel C.C., Adelberger K.L., Giavalisco M., Dickinson M., Pettini M., 1999, ApJ (in press), astro-ph/9811399
 Treyer M.A., Ellis R.S., Milliard B., Donas J., Bridges T.J., 1998, MN 300, 303
 Walker T.P., Steigman G., Schramm D.N., Olive K.A., Kang H.-S., 1991, ApJ 376, 51
 Williams R.E. et al, 1996, AJ 112, 1335